# DAFTAR PUSTAKA


Alves, R. et al. (2013). Antifungal Constituents from the Roots of Piper dilatatum Rich . , 2013.

Ang, M. (2014). Antibacterial activity of the Piper aduncum oil and dillapiole , its main constituent , against multidrug-resistant strains, *13*(6), 517–526

Atun, Sri. (2014). Metode Isolasi Dan Identifikasi Struktur Senyawa Organik Bahan Alam.8(2):53-61

Bangash, F.A. et al., (2012). In-Vitro Antibacterial Activity Of Piper Betel Leaf Extracts. , 03(04), pp.639–646.

Campos, M. P., Cechinel, V., Silva, R. Z., & Yunes, R. A. (2007). Antibacterial Activity of Extract , Fractions and Four Compounds Extracted from Piper solmsianum C . DC . var . solmsianum ( Piperaceae ).

Darwis, D. (2000). Teknik Dasar Laboratorium Dalam Penelitian Senyawa Bahan Alam Hayati. Workshop Pengembangan Sumber Daya Manusia Dalam Bidang Kimia Organik Bahan Alam Hayati. FMIPA Universitas Andalas Padang

Dewick, P. M. (2009). *Medicinal Natural Products–A Biosynthetic Approach* (3rd ed., pp. 21, 147-153, 166-169,326-328). United Kingdom: John Wiley & Sons Ltd.

Dinanti, B. R. (2014). Long Pepper ( Piper retrofractum Vahl ) to Overcome Erectile Dysfunction, *3*, 3–8.

Evizal, R. (1996). Pembibitan cabe jawa menggunakan setek pendek. Seri Monografi LP Unila (4):68- 78.

Evizal, R. (2013). Tanaman Rempah dan Fitofarmaka. Fakultas Pertanian Unila. Bandar Lampung.

Hardik, S.B. et al. (2007). Antileishmanial amides and lignans from *Piper cubeba* and *Piper retrofractum.* J Nat Med;61:418-421.

Hendayana, Sumar. (1994). Kimia Analitik Instrumen. Semarang : Semarang Press.

James, T.L. (1998). Fundamental of NMR. Dept of Pharmaceutical Chemistry Univ. Of California

Jensen S, Hansen J, Boll PM. (1993). Lignans and neolignans from *Piperaceae*.

Phytochemistry; 33:523-530.

Kelly, J. et al. (2014). Antifungal Activity and Computational Study of Constituents from Piper divaricatum Essential Oil against Fusarium Infection in Black Pepper. , pp.17926–17942.

Kristanti, A.N., Aminah, N.S., Tanjung, M., dan Kurniadi, B. (2008). Buku Ajar FITOKIMIA. Surabaya : Airlangga University Press.

Lee, S.A. et al. (2008). Methyl Piperate Derivatives from Piper longum and Their Inhibition of Monoamine Oxidase. , 31(6), pp.679–683.

Lie CL, Chien CS, Yuh CS, Tung HT. (2006). Anti-inflammatory neolignans from *Piper kadsura.* J Nat Prod ; 69:842-844.

Mohd, W. et al.(2014). Anticholinesterase and Antityrosinase Activities of Ten Piper Species from Malaysia. , 4(Suppl 2), pp.527–531.

Morikawa, T. et al. (2004). New Amides and Gastroprotective Constituents from the Fruit of Piper chaba.

Nakatani, N., Inatani, R., Ohta, H., & Nishioka, A. (1986). Chemical Constituents of Peppers ( Piper spp .) and Application to Food Preservation : Naturally Occurring Antioxidative Compounds, *67*, 135–142.

Okwute, S.K. et al. (2013). Piperine-Type Amides : Review of the Chemical and Biological Characteristics. , 5(3), pp.99–122.

Salleh WMNHW, Ahmad F, Yen KH. (2014). Chemical composition and antimicrobial activity of essential oil of *Piper muricatum* Blume (Piperaceae) J Essent Oil Bear Pl ; 17:1329-1334.

Sengupta S, Ryan AB. (1987). The chemistry of *Piper* species: a review. Fitoterapia ; 58:147-165.

Siddiqui BS, Gulzar T, Mahmood A, Begum S, Khan B, Afshan F. (2004). New insecticidal amides from petroleum ether extract of dried *Piper nigrum* L. whole fruits. Planta Med ; 52:1349-1352.

Tabudravu, J. N., and Jaspars, M. (2005). Anticancer activities of constituents of kava (*Piper methysticum*) : *The South Pacific Journal of Natural Science,* Vol. 23.

Wu QL, Wang SP, Tu GZ, Feng YX, Yang JS. (1997). Alkaloids from *Piper puberullum*. Phytochemistry ; 44:727-730.

Yamaguchi L, Lago JHG, Tanizaki TM, Mascio PD, Kato MJ. (2006). Antioxidant activity of prenylated hydroquinone and benzoic acid derivatives from *Piper crassinervium* Kunth. Phytochemistry ;67:1838-184.